PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau 1.44



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT) (51) International Patent Classification 6: WO 99/10514 (11) International Publication Number: A1 C12N 15/82, 15/31, A01H 5/00 (43) International Publication Date: 4 March 1999 (04.03.99) PCT/US98/17546 (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, (21) International Application Number: BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, (22) International Filing Date: 25 August 1998 (25.08.98) LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian (30) Priority Data: 60/057,562 26 August 1997 (26.08.97) US patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, (71) Applicant (for all designated States except US): NORTH IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). CAROLINA STATE UNIVERSITY [US/US]; 1 Holladay Hall, Campus Box 7003, Raleigh, NC 27695-7003 (US). Published (72) Inventors; and (75) Inventors/Applicants (for US only): OBEID, Lina, M. With international search report. [US/US]; 104 Crowther Court, Chapel Hill, NC 27514 (US). BOSS, Wendy, F. [US/US]; 8621 Carswell Court, Raleigh, NC 27613 (US). MAO, Cungui [CN/US]; 941 Lambeth Circle, 18B, Durham, NC 27705 (US). (74) Agents: BENNETT, Virginia, C. et al.; Myers, Bigel, Sibley, & Sajovec, P.A., P.O. Box 37428, Raleigh, NC 27627 (US). (54) Title: FUMONOSIN RESISTANCE

(57) Abstract

DNA encoding a protein capable of increasing resistance to mycotoxins of the fumonosin family is described, as are gene transfer vectors useful for imparting fumonosin resistance (e.g., resistance to fumonosin B1) to a plant or animal. The vector comprises an expression cassette, the expression cassette contains a DNA encoding a fumonosin-resistance protein. Methods of making fumonosin-resistant transgenic plants and animals, and fumonosin-resistant transgenic plants and animals, are also described.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
Cĭ	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

FUMONOSIN RESISTANCE

This application claims the benefit of U.S. Provisional Application No. 60/057,562; filed 26 August 1997.

This invention was made with Government support under grant number IR29-AG-12467 from the National Institutes of Health. The Government has certain rights to this invention.

Field of the Invention

The present invention relates to DNA encoding proteins that increase resistance to the mycotoxin fumonosin in plant and animal cells, and to transgenic plants and animals having increased resistance to mycotoxins of the fumonosin family, such as fumonosin B1.

Background of the Invention

The fumonosins are a family of mycotoxins that are common contaminants of maize, sorghum and related grains throughout the world. These compounds were first identified in a study of a high incidence of oesophageal cancer in certain villagers in the Transkei region of South Africa. The villagers consumed beer brewed from moldy corn infected by *Fusarium moniliforme*, which produces fumonosin B1. A. Merrill et al., *Trends in Cell Biology* 6, 218 (June 1996).

15

20

Fumonosin B1, the most common of the fumonosins, is produced by F. *moniliforme* (Sheldon). The toxin is also implicated in two devastating and costly diseases of veterinary animals: equine leukoencephalomacin and porcine pulmonary oedema. Id. Fumonosin B1 is both toxic and carcinogenic to plants and animals.

In view of the foregoing, it would be extremely useful to have a means for imparting fumonosin resistance to plants, particularly grains and other monocots, as well as plants and animals susceptible to infection with a fumonosin-producing fungi.

10

20

25

30

Summary of the Invention

A first aspect of the present invention is a plant or animal gene transfer vector useful for imparting fumonosin resistance to a plant or animal. The vector comprising an expression cassette, the expression cassette contains a DNA encoding a fumonosin-resistance protein (e.g., an ATP-binding cassette transporter). In general, such a DNA is (a) a DNA having a sequence according to SEQ ID NO:1, (b) a DNA that hybridizes to DNA having a sequence according to SEQ ID NO:1 and encodes and an ATP-binding cassette transporter that imparts fumonosin-resistance to a plant or animal cell, or (c) a DNA that encodes a protein encoded by a DNA of (a) or (b) above, but differs from the DNA of (a) or (b) above due to the degeneracy of the genetic code.

A second aspect of the present invention is a method of making a fumonosin-resistant transgenic plant. The method comprises transforming a plant cell with an expression cassette as described above, and then regenerating a fumonosin-resistant transgenic plant from the transformed plant cell.

A third aspect of the present invention is a fumonosin-resistant transgenic plant, wherein some or all of the cells of the plant contain a heterologous expression cassette as described above.

A fourth aspect of the present invention is a method of making a fumonosin-resistant transgenic animal. The method comprises transforming an

animal cell with an expression cassette as described above, and then regenerating a fumonosin-resistant transgenic animal from the transformed animal cell.

A fifth aspect of the present invention is a fumonosin-resistant transgenic non-human animal, wherein some or all of the cells of the animal containing a heterologous expression cassette as described above.

The foregoing and other objects and aspects of the present invention are explained in greater detail in the specification set forth below.

Detailed Description of the Invention

The present invention may be used to impart resistance to any type of fumonosin to plants and animals, including fumonosins of the A, B, and C series (e.g., fumonosin A1, fumonosin B1, fumonosin B2, fumonsin C1, phytotoxin TA). The imparting of resistance to B series fumonosins is preferred, and the imparting of resistance to fumonosin B1 is most preferred. The term "resistance" as used herein does not imply complete resistance, but rather refers to any increase in the level of resistance that is of a commercial agriculatural or veterinary advantage as compared to the same animal without the presence of the expression cassette.

A. DNA sequences

10

15

20

25

30

DNAs sequences useful for carrying out the present invention include those coding for ATP-binding cassette (ABC) transporters, and particularly for proteins homologous to, and having essentially the same biological properties as, the protein given herein SEQ ID NO:2. This definition is intended to encompass natural allelic variations therein. Isolated DNA or cloned genes of the present invention can be of any species of origin, including microorganism, plant, and animal, (see generally C. Higgins, ABC Transporters: from microorganisms to man, Annu. Rev. Cell Biol. 8: 67 (1992)), but are typically of natural origin and are preferably of yeast origin. Thus, DNAs which hybridize to DNA disclosed herein as SEQ ID NO:1 (or fragments or derivatives thereof which serve as hybridization probes as discussed below) and which code on expression for a protein of the present

invention (e.g., a protein according to SEQ ID NO:2) are also an aspect of this invention.

Conditions which will permit other DNAs which code on expression for a protein of the present invention to hybridize to the DNA of SEQ ID NO:1 disclosed herein can be determined in accordance with known techniques. For example, hybridization of such sequences may be carried out under conditions of reduced stringency, medium stringency or even stringent conditions (e.g., conditions represented by a wash stringency of 35-40% Formamide with 5x Denhardt's solution, 0.5% SDS and 1x SSPE at 37°C; conditions represented by a wash stringency of 40-45% Formamide with 5x Denhardt's solution, 0.5% SDS, and 1x SSPE at 42°C; and conditions represented by a wash stringency of 50% Formamide with 5x Denhardt's solution, 0.5% SDS and 1x SSPE at 42°C, respectively) to DNA of SEQ ID NO:1 disclosed herein in a standard hybridization assay. See, e.g., J. Sambrook et al., Molecular Cloning, A Laboratory Manual (2d Ed. 1989)(Cold Spring Harbor Laboratory)). In general, sequences which code for proteins of the present invention and which hybridize to the DNA of SEQ ID NO:1 disclosed herein will be at least 75% homologous, 85% homologous, and even 95% homologous or more with SEQ ID NO:1.

DNAs which code for proteins of the present invention, or DNAs which hybridize to that of SEQ ID NO:1, but which differ in codon sequence from SEQ ID NO:1 due to the degeneracy of the genetic code, are also an aspect of this invention. The degeneracy of the genetic code, which allows different nucleic acid sequences to code for the same protein or peptide, is well known in the literature. See, e.g., U.S. Patent No. 4,757,006 to Toole et al. at Col. 2, Table 1.

20

25

Knowledge of the nucleotide sequence as disclosed herein in SEQ ID NO:1 can be used to generate hybridization probes which specifically bind to the DNA of the present invention or to mRNA to determine the presence of amplification or overexpression of the proteins of the present invention. Pairs of probes which will serve as PCR primers for the DNA sequences of the present invention, or portions thereof, may be used in accordance with the process described in U.S. Patents Nos. 4,683,202 and 4,683,195 to Mullis (applicant specifically intends that the

disclosures of all U.S. Patent references disclosed herein be incorporated herein by reference).

Since numerous ATP-binding cassette (ABC) transporters are known, see, e.g., C. Higgins, Annu. Rev. Cell Biol. 8, 67 (1992), ABC transporters that impart fumonosin resistance to plant or animal cells when expressed therein can also be identified by expressing that transporter in a plant or animal cell, or a yeast cell, and then testing that cell for fumonosin resistance, essentially as described below.

B. Genetic Engineering Techniques

5

10

20

25

30

The production of cloned genes, recombinant DNA, vectors, transformed host cells, proteins and protein fragments by genetic engineering is well known. *See*, e.g., U.S. Patent No. 4,761,371 to Bell et al. at Col. 6 line 3 to Col. 9 line 65; U.S. Patent No. 4,877,729 to Clark et al. at Col. 4 line 38 to Col. 7 line 6; U.S. Patent No. 4,912,038 to Schilling at Col. 3 line 26 to Col. 14 line 12; and U.S. Patent No. 4,879,224 to Wallner at Col. 6 line 8 to Col. 8 line 59. (Applicant specifically intends that the disclosure of all U.S. patent references cited herein be incorporated by reference herein in their entirety).

DNA constructs of the present invention may be used to transform cells from a variety of organisms, including plants (*i.e.*, vascular plants) and animals (particularly mammals such as horses, cows and pigs). As used herein, plants includes both gymnosperms and angiosperms (*i.e.*, monocots and dicots). Transformation according to the present invention may be used to increase expression levels of transgenes in stably transformed cells.

The term "operatively associated," as used herein, refers to DNA sequences on a single DNA molecule which are associated so that the function of one is affected by the other. Thus, a transcription initiation region is operatively associated with a structural gene when it is capable of affecting the expression of that structural gene (i.e., the structural gene is under the transcriptional control of the transcription initiation region). The transcription initiation region is said to be "upstream" from the structural gene, which is in turn said to be "downstream" from the transcription initiation region.

10

20

25

30

DNA constructs, or "expression cassettes," of the present invention preferably include, 5' to 3' in the direction of transcription, a transcription initiation region, a structural gene operatively associated with the transcription initiation region, and a termination sequence including a stop signal for RNA polymerase and a polyadenylation signal for polyadenylation (e.g., the nos terminator. All of these regions should be capable of operating in the cells to be transformed. Matrix attachment regions flanking the expression cassette may optionally be included. The termination region may be derived from the same gene as the transcription initiation or promoter region, or may be derived from a different gene.

The transcription initiation region, which preferably includes the RNA polymerase binding site (promoter), may be native to the host organism to be transformed or may be derived from an alternative source, where the region is functional in the host. Other sources include the Agrobacterium T-DNA genes, such as the transcriptional initiation regions for the biosynthesis of nopaline, octapine, mannopine, or other opine transcriptional initiation regions, transcriptional initiation regions from plants, transcriptional initiation regions from viruses (including host specific viruses), or partially or wholly synthetic transcription initiation regions. Transcriptional initiation and termination regions are well known. *See, e.g.*, dGreve, *J. Mol. Appl. Genet.* 1, 499-511 (1983); Salomon et al., *EMBO J.* 3, 141-146 (1984); Garfinkel et al., *Cell* 27, 143-153 (1983); and Barker et al., *Plant Mol. Biol.* 2, 235-350 (1983).

The transcriptional initiation regions may, in addition to the RNA polymerase binding site, include regions which regulate transcription, where the regulation involves, for example, chemical or physical repression or induction (e.g., regulation based on metabolites or light) or regulation based on cell differentiation (such as associated with leaves, roots, seed, or the like in plants). Thus, the transcriptional initiation region, or the regulatory portion of such region, is obtained from an appropriate gene which is so regulated. For example, the 1,5-ribulose biphosphate carboxylase gene is light-induced and may be used for transcriptional initiation. Other genes are known which are induced by stress, temperature, wounding, pathogen effects, etc. Tissue specific promoters, such as

10

15

20

25

30

root-specific promoters or a promoter specific for corn silk, may advantageously be employed as will be apparent to those skilled in the art. In corn for the prevention of corn ear rot (or "pink ear rot of maize"), a promoter associated with Pedicel Glutamine Synthetase gene (preferentially expressed in the region of the kernel attached to the cob, where fumonosin-fungi enter the kernel) may be employed, or a promoter that preferentially expresses in the seed or seed coat.

The expression cassette may be provided in a DNA construct that also has at least one replication system. For convenience, it is common to have a replication system functional in Escherichia coli, such as ColE1, pSC101, pACYC184, or the like. In this manner, at each stage after each manipulation, the resulting construct may be cloned, sequenced, and the correctness of the manipulation determined. In addition, or in place of the E. coli replication system, a broad host range replication system may be employed, such as the replication systems of the P-1 incompatibility plasmids, e.g., pRK290. In addition to the replication system, there will frequently be at least one marker present, which may be useful in one or more hosts, or different markers for individual hosts. That is, one marker may be employed for selection in a prokaryotic host, while another marker may be employed for selection in a eukaryotic host, particularly a plant host. The markers may be protection against a biocide, such as antibiotics, toxins, heavy metals, or the like; provide complementation, for example by imparting prototrophy to an auxotrophic host; or provide a visible phenotype through the production of a novel compound. Exemplary genes that may be employed include neomycin phosphotransferase (NPTII), hygromycin phosphotransferase (HPT), chloramphenicol acetyltransferase (CAT), nitrilase, and the gentamicin resistance gene. For plant host selection, nonlimiting examples of suitable markers are β-glucuronidase, providing indigo production, luciferase, providing visible light production, NPTII, providing kanamycin resistance or G418 resistance, HPT, providing hygromycin resistance, and the mutated aroA gene, providing glyphosate resistance.

The various fragments comprising the various constructs, expression cassettes, markers, and the like may be introduced consecutively by restriction enzyme cleavage of an appropriate replication system, and insertion of the

particular construct or fragment into the available site. After ligation and cloning the DNA construct may be isolated for further manipulation. All of these techniques are amply exemplified in the literature and find particular exemplification in Sambrook et al., Molecular Cloning: A Laboratory Manual, (2d Ed. 1989)(Cold Spring Harbor Laboratory, Cold Spring Harbor, NY).

C. Plant Genetic Engineering

10

15

20

30

As noted above, the present invention provides a method of making a fumonosin-resistant transgenic plant. The term "plant" as used herein refers to vascular plants (e.g., gymnosperms and angiosperms). The method comprises transforming a plant cell with an expression cassette as described above, and then regenerating a fumonosin-resistant transgenic plant from the transformed plant cell. The transforming step may be carried out by any suitable means, including by Agrobacterium-mediated transformation and non-Agrobacterium-mediated transformation, as discussed in detail below. Plants are regenerated from the transformed cell (or cells) by techniques known to those skilled in the art, as also discussed below. Where chimeric plants are produced by the process, plants in which all cells are transformed may be regenerated from chimeric plants having transformed germ cells, as is known in the art.

Vectors that may be used to transform plant tissue with DNA constructs/expression cassettes of the present invention include both *Agrobacterium* and non-*Agrobacterium* vectors, particularly ballistic vectors, as well as vectors suitable for DNA-mediated transformation.

Agrobacterium mediated transformation. Agrobacterium-mediated gene transfer exploits the natural ability of Agrobacterium tumefaciens to transfer DNA into plant chromosomes. Agrobacterium is a plant pathogen that transfers a set of genes encoded in a region called T-DNA of the Ti plasmid into plant cells at wound sites. The typical result of gene transfer is a tumorous growth called a crown gall in which the T-DNA is stably integrated into a host chromosome. The ability to cause crown gall disease can be removed by deletion of the genes in the T-DNA without loss of DNA transfer and integration. The DNA to be

10

15

20

25

30

transferred is attached to border sequences that define the end points of an integrated T-DNA.

The Agrobacterium strain utilized in the methods of the present invention is modified to contain a gene or genes of interest, or a nucleic acid to be expressed in the transformed cells. The nucleic acid to be transferred is incorporated into the T-region and is flanked by T-DNA border sequences. A variety of Agrobacterium species are known in the art particularly for dicotyledon transformation. Such Agrobacterium can be used in the methods of the invention. See, e.g., Hooykaas, Plant Mol. Biol. 13, 327 (1989); Smith et al., Crop Science 35, 301 (1995); Chilton, Proc. Natl. Acad. Sci. USA 90, 3119 (1993); Mollony et al., Monograph Theor. Appl. Genet NY 19, 148 (1993); Ishida et al., Nature Biotechnol. 14, 745 (1996); and Komari et al., The Plant Journal 10, 165 (1996), the disclosures of which are incorporated herein by reference.

In addition to the T-region, the Ti plasmid contains a vir region. The vir region is important for efficient transformation, and appears to be species-specific. Binary vector systems have been developed where the manipulated disarmed T-DNA carrying foreign DNA and the vir functions are present on separate plasmids. In this manner, a modified T-DNA region comprising foreign DNA (the nucleic acid to be transferred) is constructed in a small plasmid which replicates in E. coli. This plasmid is transferred conjugatively in a tri-parental mating into Agrobacterium tumefaciens that contains a compatible plasmid-carrying virulence gene. The vir functions are supplied in trans to transfer the T-DNA into the plant genome. Such binary vectors are useful in the practice of the present invention.

Preferred vectors of the present invention are super-binary vectors. See, e.g., United States Patent No. 5,591,615 and EP 0 604 662. Such a super-binary vector has been constructed containing a DNA region originating from the virulence region of the Ti plasmid pTiBo542 (Jin et al., J. Bacteriol. 169, 4417 (1987)) contained in a super-virulent Agrobacterium tumefaciens A281 exhibiting extremely high transformation efficiency (Hood et al., Biotechnol. 2, 702 (1984);

10

15

20

25

30

Hood et al., J. Bacteriol. 168, 1283 (1986); Komari et al., J. Bacteriol. 166, 88 (1986); Jin et al., J. Bacteriol. 169, 4417 (1987); Komari, Plant Science 60, 223 (1987); ATCC Accession No. 37394. Exemplary super-binary vectors known to those skilled in the art include pTOK162 (Japanese patent Appl. (Kokai) No. 4-222527, EP 504,869, EP 604,662, and United States Patent No. 5,591,616, herein incorporated by reference) and pTOK233 (Komari, Plant Cell Reports 9.303 (1990); Ishida et al., Nature Biotechnology 14, 745 (1996); herein incorporated by reference). Other super-binary vectors may be constructed by the methods set forth in the above references. Super-binary vector pTOK162 is capable of replication in both E. coli and in A. tumefaciens. Additionally, the vector contains the virB, virC and virG genes from the virulence region of pTiBo542. The plasmid also contains an antibiotic resistance gene, a selectable marker gene, and the nucleic acid of interest to be transformed into the plant. The nucleic acid to be inserted into the sorghum genome is located between the two border sequences of the T region. Super-binary vectors of the invention can be constructed having the features described above for pTOK162. The T-region of the super-binary vectors and other vectors for use in the invention are constructed to have restriction sites for the insertion of the genes to be delivered. Alternatively, the DNA to be transformed can be inserted in the T-DNA region of the vector by utilizing in vivo homologous recombination. See, Herrera-Esterella et al., EMBO J. 2, 987 (1983); Horch et al., Science 223, 496 (1984). Such homologous recombination relies on the fact that the super-binary vector has a region homologous with a region of pBR322 or other similar plasmids. Thus, when the two plasmids are brought together, a desired gene is inserted into the super-binary vector by genetic recombination via the homologous regions.

Non-Agrobacterium mediated transformation. Microparticles carrying a DNA construct of the present invention, which microparticles are suitable for the ballistic transformation of a cell, are also useful as a vector for transforming cells according to the present invention. The microparticle is propelled into a cell to produce a transformed cell. Where the transformed cell is a plant cell, a plant may be regenerated from the transformed cell according to techniques known in the art.

Any suitable ballistic cell transformation methodology and apparatus can be used in practicing the present invention. Exemplary apparatus and procedures are disclosed in Sanford and Wolf, U.S. Patent No. 4,945,050. When using ballistic transformation procedures, the expression cassette may be incorporated into a plasmid capable of replicating in the cell to be transformed. Examples of microparticles suitable for use in such systems include 1 to 5 μ m gold spheres. The DNA construct may be deposited on the microparticle by any suitable technique, such as by precipitation.

Plant species may be transformed with the DNA construct of the present invention by the DNA-mediated transformation of plant cell protoplasts and subsequent regeneration of the plant from the transformed protoplasts in accordance with procedures well known in the art.

10

15

20

25

30

Any plant tissue capable of subsequent clonal propagation, whether by organogenesis or embryogenesis, may be transformed with a vector of the present invention. The term "organogenesis," as used herein, means a process by which shoots and roots are developed sequentially from meristematic centers; the term "embryogenesis," as used herein, means a process by which shoots and roots develop together in a concerted fashion (not sequentially), whether from somatic cells or gametes. The particular tissue chosen will vary depending on the clonal propagation systems available for, and best suited to, the particular species being transformed. Exemplary tissue targets include leaf disks, pollen, embryos, cotyledons, hypocotyls, megagametophytes, callus tissue, existing meristematic tissue (e.g., apical meristems, axillary buds, and root meristems), and induced meristem tissue (e.g., cotyledon meristem and hypocotyl meristem).

Plants of the present invention may take a variety of forms. The plants may be chimeras of transformed cells and non-transformed cells; the plants may be clonal transformants (e.g., all cells transformed to contain the expression cassette); the plants may comprise grafts of transformed and untransformed tissues (e.g., a transformed root stock grafted to an untransformed scion in citrus species). The transformed plants may be propagated by a variety of means, such as by clonal propagation or classical breeding techniques. For example, first generation (or T1)

transformed plants may be selfed to give homozygous second generation (or T2) transformed plants, and the T2 plants further propagated through classical breeding techniques. A dominant selectable marker (such as *npt* II) can be associated with the expression cassette to assist in breeding.

Plants that may be employed in practicing the present invention include (but are not limited to) maize or corn (Zea mays), sorghum, wheat, oats, rye, barley, rice, tobacco (Nicotiana tabacum), potato (Solanum tuberosum), soybean (glycine max), peanuts (Arachis hypogaea), cotton (Gossypium hirsutum), sweet potato (Ipomoea batatus), cassava (Manihot esculenta), coffee (Cofea spp.), coconut (Cocos nucifera), pineapple (Ananas comosus), citrus trees (Citrus spp.), cocoa (Theobroma cacao), tea (Camellia sinensis), banana (Musa spp.), avocado (Persea americana), fig (Ficus casica), guava (Psidium guajava), mango (Mangifera indica), olive (Olea europaea), papaya (Carica papaya), cashew (Anacardium occidentale), macadamia (Macadamia integrifolia), almond (Prunus amygdalus), sugar beets (Beta vulgaris), vegetables, ornamentals, and conifers.

Particularly preferred plants for carrying out the present invention are maize (Zea mays) and sorghum.

D. Transgenic Animals.

5

10

15

20

25

30

A method of making a fumonosin-resistant transgenic animal is also an aspect of the present invention. The method can be carried out on any suitable animal subject, but is preferably carried out with non-human mammals. Ovine, bovine, and equine species are particularly preferred (e.g., pigs, cows, and horses).

The method comprises transforming an animal cell with an expression cassette as described above, in an animal transformation vector, and then regenerating a fumonosin-resistant transgenic animal from the transformed animal cell. The transformation step may be carried out by any suitable means, as discussed in detail below, and the regeneration step may also be carried out by any suitable means, as also discussed in detail below. Where chimeric animals are produced by the process, animals in which all cells are transformed may be

regenerated from chimeric animals having transformed germ cells, as is known in the art.

The production of transgenic animals can be carried out by any suitable technique, such as pronuclear microinjection, infection of embryos with retroviruses, embryonic stem cell-mediated techniques, transfer of entire chromosomal segments and gamete transfection in conjunction with *in vitro* fertilization, etc. See generally Charles River Laboratories, Transgenic Animal Science: Principles and Methods (Summer 1991).

Transgenic animals that express an ABC transporter protein can be produced by the genetic transformation of zygotes, as described in T. Wagner et al., U.S. Patent No. 4,873,191 (applicant intends that the disclosure of all U.S. Patent References cited herein be incorporated herein by reference).

10

15

20

Methods of producing a transgenic bovine or transgenic bovine embryo are described in U.S. Patent No. 5,663,076 to H. DeBoer et al.

In another technique, a pluripotent embryonic stem cell from the species to be transformed may be derived, the expression cassette inserted into the stem cell, and one or more of the stem cells inserted into an early embryo such as a blastocyst of the animal to be transformed, and the animal raised to birth in a suitable female host (e.g., M. Evans, PCT Application WO90/03432).

In still another technique, embryonic stem cells useful for making chimeric and transgenic ungulates (e.g., porcine, bovine, ovine and caprine species) are described in M. Wheeler, PCT Application WO 94/26884. In general, the embryonic stem cells are transformed with the exogenous genetic material of interest (e.g., an ABC transporter expression cassette) and then used to provide chimeric ungulates which have germ cells comprising the exogenous genetic material. The chimeric ungulates are bred to provide transgenic ungulates (see also U.S. Patent No. 5,523,226 to M. Wheeler).

Methods of producing transgenic animals by subjecting a mixture of DNA and the embryo to an electric discharge are described in U.S. Patent No. 5,567,607 to X. Zhao et al.

Mammalian expression vectors are described in U.S. Patent No. 5,627,033 to J. Smith et al.

E. Utilities

The methods, constructs, and products described above are useful in providing a selectable marker (i.e., Fumonosin resistance) for genetic engineering techniques, where other nucleic acid segments are being introduced into the plant or animal cell in association with the fumonosin-resistance gene.

The methods, constructs, and products described above are useful in providing plants and animals that are resistant, or have greater levels of resistance, to naturally occurring fumonosin infection.

The examples that follow are provided to illustrate the present invention, and are not to be construed as limiting thereof.

15

20

25

10

5

EXPERIMENTAL

A Fumonosin B1 sensitive Sacharomyces cerevisiae strain (JS16) was screened, which was sensitive to 400 μM Fumonosin B1 in synthetic complete medium at a cell density as high as 10⁷ cells/mL (Wild-type yeast strain JK93da is resistant to as high as 1 mM Fumonosin B1). Then, a S. cerivisiae genomic DNA library was constructed in the multicopy vector YEP24. JS16 was transformed with the DNA library and four Fumonosin B1 resistant clones were selected on synthetic complete agar medium with 400 μM Fumonosin B1.

DNA sequencing was carried out among the four clones. Two of them had the same DNA sequence, the other two had the consensus DNA sequences as the first two clones. The consensus sequence (SEQ ID NO:1) encodes a protein of 1477 amino acid residues (SEQ ID NO:2). Overexpression of the protein confers Fumonosin B1 resistance to S. cerivasiae.

The gene deletion mutant is sensitive to Fumonosin B1 compared to wild type strain (JK93da).

A yeast database was searched. It was found that the protein belongs to an ABC transport family. See D. Katzmann et al., Molecular and Cellular Biology 15, 6875 (1995). The gene is located on chromosome VII.

That which is claimed is:

10

1. A DNA construct comprising an expression cassette, which construct comprises, in the 5' to 3' direction, a promoter operable in a plant cell and a DNA segment according to claim 1 positioned downstream from said promoter and operatively associated therewith, said DNA segment selected from the group consisting of:

- (a) SEQ ID NO:1;
- (b) DNA sequences which encode an enzyme having SEQ ID NO:2:
- (c) DNA sequences which hybridize to isolated DNA of (a) or(b) above and which encode an ATP-binding cassette transporter; and(d) DNA sequences which differ from the DNA of (a), (b) or(c) above due to the degeneracy of the genetic code.
 - 2. A DNA construct according to claim 1, wherein said promoter is constitutively active in plant cells.
 - 3. A DNA construct according to claim 1, wherein said promoter is selectively active in plant tissue cells.
 - 4. A DNA construct according to claim 3, wherein said promoter is selectively active in plant cells selected from root cells, seed cells, seed coat cells, and corn silk cells.
 - 5. A DNA construct according to claim 3, wherein said promoter is a corn Pedicel Glutamine Synthetase promoter that is expressed in the region of the corn kernel attached to the cob.

6. A DNA construct according to claim 1, wherein said construct further comprises a plasmid.

- 7. A DNA construct according to claim 1 carried by a plant transformation vector.
- 8. A DNA construct according to claim 1 carried by a plant transformation vector, which plant transformation vector is an *Agrobacterium tumefaciens* vector.
 - 9. A plant cell containing a DNA construct according to claim 1.
 - 10. A transgenic plant comprising plant cells according to claim 9.
- 11. A method of making a transgenic plant cell having increased resistance to fumonosin, said method comprising:

providing a plant cell;

providing an exogenous DNA construct, which construct comprises, in the 5' to 3' direction, a promoter operable in a plant cell and a DNA sequence that encodes an ATP-binding cassette transporter protein that increases resistance of a plant cell to fumonosin, said DNA sequence operably associated with said promoter; and

transforming said plant cell with said DNA construct to produce a
transformed plant cell, said plant cell having increased resistance to fumonosin
compared to an untransformed cell.

12. The method of claim 11, wherein said plant cell is a corn cell.

13. The method of claim 11, further comprising regenerating a plant from said transformed plant cell.

- 14. A method according to claim 11, wherein said promoter is constitutively active.
- 15. A method according to claim 11, wherein said promoter is selectively active in plant cells.
- 16. A method according to claim 11, wherein said promoter is selectively active in plant cells selected from root cells, seed cells, seed coat cells, and corn silk cells.
- 17. A method according to claim 11, wherein said promoter is a corn Pedicel Glutamine Synthetase promoter that is expressed in the region of the corn kernel attached to the cob.
- 18. A method according to claim 11, wherein said transforming step is carried out by bombarding said plant cell with microparticles carrying said DNA construct.
- 19. A method according to claim 11 wherein said transforming step is carried out by infecting said plant cell with an *Agrobacterium tumefaciens* containing a Ti plasmid carrying said DNA construct.
- 20. A method of producing transgenic plant seeds, comprising collecting seed from a transgenic plant produced by the method of claim 11.
- 21. The method according to claim 11, wherein said exogenous DNA sequence comprises a DNA sequence selected from the DNA sequences of Claim 1.

22. A transgenic plant having increased resistance to fumonosin compared to a non-transformed control plant, said transgenic plant comprising transgenic plant cells containing:

an exogenous DNA construct comprising, in the 5' to 3' direction, a promoter operable in said plant cell and a DNA sequence that encodes an ATP-binding cassette transporter protein that increases resistance of a plant cell to fumonosin, said DNA sequence operably associated with said promoter;

said plant exhibiting increased fumonosin resistance compared to a non-transformed control plant.

5

- 23. The method according to claim 22, wherein said exogenous DNA sequence comprises a DNA sequence selected from the DNA sequences of claim 1.
- 24. A plant according to claim 22, wherein said promoter is a constitutively active promoter.
- 25. A plant according to claim 22, wherein said promoter is selectively active in plant cells.
- 26. A method according to claim 22, wherein said promoter is selectively active in plant cells selected from root cells, seed cells, seed coat cells, and corn silk cells.
- 27. A method according to claim 22, wherein said promoter is a corn Pedicel Glutamine Synthetase promoter that is expressed in the region of the corn kernel attached to the cob.
 - 28. A transgenic plant according to claim 22, which plant is corn.

29. A transgenic corn plant having increased fumonosin resistance compared to a non-transformed control plant, wherein said transgenic plant is a progeny of a plant according to claim 28.

- 30. Seeds of a transgenic corn plant having increased fumonosin resistance relative to a non-transformed control plant, wherein said transgenic plant is a plant according to claim 28 or a progeny thereof.
- 31. A method of producing transgenic plant seeds, comprising collecting seed from a transgenic plant produced by the method of claim 22.

1 SEQUENCE LISTING

<110> Obeid, Lina M.

Boss, Wendy F.

Mao, Cungui

<120> Fumonosin Resistance Proteins

<130> Obeid et al.

<140>

<141>

<150> 60/057,562

<151> 1997-08-26

<160> 2

<170> PatentIn Ver. 2.0

<210> 1

<211> 4434

<212> DNA

<213> Saccharomyces cerevisiae

<220>

<221> CDS

<222> (1)..(4434)

VO 99/10514	·	PCT/US98/17546

									2							
<400	0> 1															•
atg	acg	att	acc	gtg.	9 99	gat	gca	gtt	tcg	gag	acg	gag	ctg	gaa	aac	48
Met	Thr	Ile	Thr	Val	Gly	Asp	Ala	Val	Ser	Glu	Thr	Glu	Leu	Glu	Asn	
1				5					10					15		
aaa	agt	caa	aac	gtg	gta	cta	tct	ccc	aag	gça	tct	gct	tct	tca	gac	96
Lys	Ser	Gln	Asn	Val	Val	Leu	Ser	Pro	Lys	Ala	Ser	Ala	Ser	Ser	Asp	
			20					25					30			
ata	agc	aca	gat	gtt	gat	aag	gac	aca	tcg	tct	tct	tgg	gat	gac	aaa	144
Ile	Ser	Thr	Asp	Val	Asp	Lys	Asp	Thr	Ser	Ser	Ser	Trp	Asp	Asp	Lys	
		35					40					45				
				·												

tct ttg ctg cct aca ggt gaa tat att gtg gac aga aat aag ccc caa 192
Ser Leu Leu Pro Thr Gly Glu Tyr Ile Val Asp Arg Asn Lys Pro Gln
50 55 60

acc tac ttg aat agc gat gat atc gaa aaa gtg aca gaa tct gat att 240

Thr Tyr Leu Asn Ser Asp Asp Ile Glu Lys Val Thr Glu Ser Asp Ile

65 70 75 80 ---

ttc cct cag aaa cgt ctg ttt tca ttc ttg cac tct aag aaa att cca 288

Phe Pro Gln Lys Arg Leu Phe Ser Phe Leu His Ser Lys Lys Ile Pro

85 90 95

gaa gta cca caa acc gat gac gag agg aag ata tat cct ctg ttc cat 336
Glu Val Pro Gln Thr Asp Asp Glu Arg Lys Ile Tyr Pro Leu Phe His
100 105 110

									3							
aca	aat	att	atc	tct	aac	atg	ttt	ttt	tgg	tgg	gtt	cta	ccc	atc	ctg	384
Thr	Asn	Ile	Ile	Ser	Asn	Met	Phe	Phe	Trp	Trp	Val	Leu	Pro	Ile	Leu	
	•	115					120					125				
cga	gtt	ggt	tat	aag	aga	acg	ata	cag	ccg	aac	gat	ctc	ttc	aaa	atg	432
Arq	Val	Gly	Tyr	Lys	Arq	Thr	Ile	Gln	Pro	Asn	Asp	Leu	Phe	Lys	Met	
_	130	-	-	-	_	135					140			-		
			_ 4	. _ .												400
															aac	480
	Pro	Arg	Met	Ser		Glu	Thr	Leu	Tyr			Phe	GIu	гла		
145					150					155					160	
atg	att	tac	tat	ttt	gag	aag	acg	agg	aaa	aaa	tac	cgt	aaa	aga	cat	528
Иet	Ile	Tyr	Tyr	Phe	Glu	Lys	Thr	Arg	Lys	Lys	Tyr	Arg	Lys	Arg	His	
				165					170					175		
cca	gaa	gcg	aca	gaa	gaa	gag	gtt	atg	gaa	aat	gcc	aaa	cta	cct	aaa	576
?ro	Glu	Ala	Thr	Glu	Glu	Glu	Val	Met	Glu	Asn	Ala	Lys	Leu	Pro	Lys	
			180					185					190			
at	aca	gtt	ctg	aga	gct	tta	tta	ttc	act	ttt	aag	aaa	cag	tac	ttc	624
			Leu													
		195					200				-1-	205		-1-		
		1,,,					200					203				
			gtg													672
let	Ser	Ile	Val	Phe	Ala	Ile	Leu	Ala	Asn	Суз	Thr	Ser	Gly	Phe	Asn	
	210					215					220					

ccc atg att acc aag agg cta att gag ttt gtc gaa gaa aag gct att 720

768

Pro	Met	Tle	Thr	T.ve	Ara	Ĩ. 2 11	alī	Glu	4 Phe	Val	Glu	Glu	Lvs	Ala	Tle
				273	••••	200		<u> </u>		•	-		_,_		
225					230					235					240
													•		
ttt	cat	agc	atg	cat	gtt	aac	aaa	ggt	att	ggt	tac	gct	att	ggt	gca
Phe	His	Ser	Met	His	Val	Asn	Lys	Gly	Ile	Gly	Tyr	Ala	Ile	Gly	Ala
				245					250					255	

tgt ttg atg atg ttc gtt aac ggg ttg acg ttc aac cat ttc ttt cat 816

Cys Leu Met Met Phe Val Asn Gly Leu Thr Phe Asn His Phe Phe His

260 265 270

aca tcc caa ctg act ggt gtg caa gct aag tct att ctt act aaa gct 864
Thr Ser Gln Leu Thr Gly Val Gln Ala Lys Ser Ile Leu Thr Lys Ala
275 280 285

gcc atg aag aaa atg ttt aat gca tct aat tat gcg aga cat tgt ttt 912
Ala Met Lys Lys Met Phe Asn Ala Ser Asn Tyr Ala Arg His Cys Phe
290 295 300

cct aac ggt aaa gtg act tct ttt gta aca aca gat ctc gct aga att ~960 Pro Asn Gly Lys Val Thr Ser Phe Val Thr Thr Asp Leu Ala Arg Ile 305 310 315 320

gaa ttt gcc tta tct ttt cag ccg ttt ttg gct ggg ttc cct gca att 1008
Glu Phe Ala Leu Ser Phe Gln Pro Phe Leu Ala Gly Phe Pro Ala Ile
325 330 335

ttg gct att tgc att gtt tta ttg atc gtt aac ctt gga ccc att gcc 1056
Leu Ala Ile Cys Ile Val Leu Leu Ile Val Asn Leu Gly Pro Ile Ala

tta gtt ggg att ggt att ttt ttc ggt ggg ttt ttc ata tcc tta ttt Leu Val Gly Ile Gly Ile Phe Phe Gly Gly Phe Phe Ile Ser Leu Phe gca ttt aag tta att ctg ggc ttt aga att gct gcg aac atc ttc act Ala Phe Lys Leu Ile Leu Gly Phe Arg Ile Ala Ala Asn Ile Phe Thr gat gct aga gtt acc atg atg aga gaa gtg ctg aat aat ata aaa atg Asp Ala Arg Val Thr Met Met Arg Glu Val Leu Asn Asn Ile Lys Met att aaa tat tat acg tgg gag gat gcg tat gaa aaa aat att caa gat Ile Lys Tyr Tyr Thr Trp Glu Asp Ala Tyr Glu Lys Asn Ile Gln Asp att agg acc aaa gag att tct aaa gtt aga aaa atg caa cta tca aga Ile Arg Thr Lys Glu Ile Ser Lys Val Arg Lys Met Gln Leu Ser Arg aat ttc ttg att gct atg gcc atg tct ttg cct agt att gct tca ttg Asn Phe Leu Ile Ala Met Ala Met Ser Leu Pro Ser Ile Ala Ser Leu gtc act ttc ctt gca atg tac aaa gtt aat aaa gga ggc agg caa cct

Val Thr Phe Leu Ala Met Tyr Lys Val Asn Lys Gly Gly Arg Gln Pro

WO 99/10514		PCT/US98/17546	
	•		

										6							
						tct											1440
	Gly	Asn	Ile	Phe	Ala	Ser	Leu	Ser	Leu	Phe	Gln	`Val	Leu	Ser	Leu	Gln	
	465					470					475					480	
	atg	ttt	ttc	tta	cct	att	gct	att	ggt	act	gga	att	gac	atg	atc	att	1488
·	Met	Phe	Phe	Leu	Pro	Ile	Ala	Ile	Gly	Thr	Gly	Ile	Asp	Met	Ile	Ile	· .
					485					490					495		
	gga	ttg	ggc	cgt	ttg	caa	agc	tta	ttg	gag	gct	cca	gaa	gat	gat	cca	1536
	Gly	Leu	Gly	Arg	Leu	Gln	Ser	Leu	Leu	Glu	Ala	Pro	Glu	Asp	Asp	Pro	
				500					505					510			
	aat	cag	atg	att	gaa	atg	aag	cct	tct	cct	ggc	ttt	gat	cca	aaa	ttg	1584
	Asn	Gln	Met	Ile	Glu	Met	Lys	Pro	Ser	Pro	Gly	Phe	Asp	Pro	Lys	Leu	,
			515					520					525				
	gct	tta	aaa	atg	aca	cat	tgc	tca	ttt	gag	tgg	gaa	gat	tat	gaa	tta	1632
	Ala	Leu	Lys	Met	Thr	His	Cys	Ser	Phe	Glu	Trp	Glu	Asp	Tyr	Glu	Leu	
		530					535					540					
	aac	gac	gct	att	gaa	gaa	gca	aaa	gga	gaa	gct	aaa	gat	gaa	ggt	aaa	1680
	Asn	Asp	Ala	Ile	Glu	Glu	Ala	Lys	Gly	Glu	Ala	Lys	Asp	Glu	Gly	Lys	
	545					550					555					560	
	aag	aac	aaa	aaa	aag	cgt	aag	gat	aca	tgg	ggt	aag	cca	tct	gca	agt	1728
	Lys	Ăsn	īуs	Ĺуs	Lys	Arg	Lys	Asp	Thr	Trp	Gly	Lys	Pro	Ser	Ala	Ser	
					565					570					575		

									7							
act	aat	aag	gcg	aaa	aga	ttg	gac	aat	atg	ttg	aaa	gac	aga	gac	ggc	1776
Thr	Asn	Lys	Ala	Lys	Arg	Leu	Asp	Asn	Met	Leu	Lys	Asp	Arg	Asp	Gly	
			580					585					590			
ccg	gaa	gat	tta	gaa	aaa	act	tcg	ttt	agg	ggt	ttc	aag	gac	ttg	aac	1824
Pro	Glu	Asp	Leu	Glu	Lys	Thr	Ser	Phe	Arg	Gly	Phe	Lys	Asp	Leu	Asn	
		595					600					605				
ttc	gat	att	aaa	aag	ggc	gaa	ttt	att	atg	att	acg	gga	cct	att	ggt	1872
Phe	Asp	Ile	Lys	Lys	Gly	Glu	Phe	Ile	Met	Ile	Thr	Gly	Pro	Ile	Gly	
	610					615					620					
act	ggt	aaa	tct	tca	tta	ttg	aat	gcg	atg	gca	gga	tca	atg	aga	aaa	1920
Thr	Gly	Lys	Ser	Ser	Leu	Leu	Asn	Ala	Met	Ala	Gly	Ser	Met	Arg	Lys	
625					630					635					640	
023																
023																
	gat	ggt	aag	gtt	gaa	gtc	aac	9 99	gac	tta	tta	atg	tgt	ggt	tat	1968
act			aag Lys													1968
act			aag Lys													1968
act				Val					Asp					Gly		1968
act Thr	Asp	Gly	Lys	Val 645	Glu	Val	Asn	Gly	Asp 650	Leu	Leu	Met	Cys	Gly 655	Tyr	1968
act Thr	Asp	Gly	Lys	Val 645 aat	Glu gca	Val	Asn gta	Gly	Asp 650 gat	Leu	Leu	Met	Cys	Gly 655 ggt	Tyr tca	
act Thr	Asp	Gly	Lys caa Gln	Val 645 aat	Glu gca	Val	Asn gta	Gly aga Arg	Asp 650 gat	Leu	Leu	Met	Cys	Gly 655 ggt	Tyr tca	
act Thr	Asp	Gly	Lys	Val 645 aat	Glu gca	Val	Asn gta	Gly	Asp 650 gat	Leu	Leu	Met	Cys ttc Phe	Gly 655 ggt	Tyr tca	
act Thr cca Pro	Asp tgg Trp	Gly att Ile	Lys caa Gln 660	Val 645 aat Asn	Glu gca Ala	val tct ser	Asn gta Val	aga Arg 665	Asp 650 gat Asp	Leu aac Asn	Leu atc Ile	Met ata Ile	ttc Phe 670	Gly 655 ggt Gly	tca Ser	
act Thr cca Pro	tgg Trp	Gly att Ile	Lys caa Gln 660	Val 645 aat Asn	gca Ala	tct Ser	gta Val gat	aga Arg 665	Asp 650 gat Asp	Leu aac Asn	Leu atc Ile	Met ata Ile	ttc Phe 670	Gly 655 ggt Gly	tca Ser	2016
act Thr cca Pro	tgg Trp	att Ile aat Asn	Lys caa Gln 660	Val 645 aat Asn	gca Ala	tct Ser	gta Val gat Asp	aga Arg 665	Asp 650 gat Asp	Leu aac Asn	Leu atc Ile	Met ata Ile gtt Val	ttc Phe 670	Gly 655 ggt Gly	tca Ser	2016
act Thr cca Pro	tgg Trp	Gly att Ile	Lys caa Gln 660	Val 645 aat Asn	gca Ala	tct Ser	gta Val gat	aga Arg 665	Asp 650 gat Asp	Leu aac Asn	Leu atc Ile	Met ata Ile	ttc Phe 670	Gly 655 ggt Gly	tca Ser	2016
act Thr cca Pro	tgg Trp ttc	att Ile aat Asn 675	Lys caa Gln 660	Val 645 aat Asn gaa Glu	gca Ala aag Lys	tct Ser tat Tyr	gta Val gat Asp	aga Arg 665 gaa Glu	Asp 650 gat Asp gta Val	aac Asn gtt Val	atc Ile cgt Arg	Met ata Ile gtt Val 685	ttc Phe 670 tgc Cys	Gly 655 ggt Gly tct ser	tca Ser ttg	2016

Lys Ala Asp Leu Asp Ile Leu Pro Ala Gly Asp Met Thr Glu Ile Gly gaa cgt ggt att act tta tct ggt ggt caa aag gca cgt atc aat tta Glu Arg Gly Ile Thr Leu Ser Gly Gly Gln Lys Ala Arg Ile Asn Leu gec agg tet gtt tat aag aag gat att tat eta tte gac gat gte Ala Arg Ser Val Tyr Lys Lys Lys Asp Ile Tyr Leu Phe Asp Asp Val cta agt gct gtc gat tct cgt gtt ggt aaa cac atc atg gat gaa tgt Leu Ser Ala Val Asp Ser Arg Val Gly Lys His Ile Met Asp Glu Cys cta acc gga atg ctt gct aat aaa acc aga att tta gca acg cat caa Leu Thr Gly Met Leu Ala Asn Lys Thr Arg Ile Leu Ala Thr His Gln ttg tca ctg att gag aga gct tct aga gtc atc gtt tta ggt act gat 2352 Leu Ser Leu Ile Glu Arg Ala Ser Arg Val Ile Val Leu Gly Thr Asp ggc caa gtc gat att ggt act gtt gat gag cta aaa gct cgt aat caa Gly Gln Val Asp Ile Gly Thr Val Asp Glu Leu Lys Ala Arg Asn Gln act ttg ata aat ctt tta caa ttc tct tct caa aat tcg gag aaa gag

Thr Leu Ile Asn Leu Leu Gln Phe Ser Ser Gln Asn Ser Glu Lys Glu

9 810

805

815

gat gaa gaa cag gaa gcg gtt gtt gcc ggt gaa ttg gga caa cta aaa 2496 Asp Glu Glu Glu Ala Val Val Ala Gly Glu Leu Gly Gln Leu Lys 820 825 830

tat gaa tca gag gta aag gaa ttg act gaa ctg aag aaa aag gct aca 2544

Tyr Glu Ser Glu Val Lys Glu Leu Thr Glu Leu Lys Lys Ala Thr

835 840 845

gaa atg tca caa act gca aat agt ggt aaa att gta gcg gat ggt cat 2592
Glu Met Ser Gln Thr Ala Asn Ser Gly Lys Ile Val Ala Asp Gly His
850 855 860

act agt agt aaa gaa gaa aga gca gtc aat agt atc agt ctg aaa ata 2640
Thr Ser Ser Lys Glu Glu Arg Ala Val Asn Ser Ile Ser Leu Lys Ile
865 870 880

tac cgt gaa tac att aaa gct gca gta ggt aag tgg ggt ttt atc gca 2688

Tyr Arg Glu Tyr Ile Lys Ala Ala Val Gly Lys Trp Gly Phe Ile Ala

885 890 895

cta ccg ttg tat gca att tta gtc gtt gga acc aca ttc tgc tca ctt 2736

Leu Pro Leu Tyr Ala Ile Leu Val Val Gly Thr Thr Phe Cys Ser Leu
900 905 910

Phe Ser Ser Val Trp Leu Ser Tyr Trp Thr Glu Asn Lys Phe Lys Asn
915
920
925

WO 99/10514	PCT/US98/17546

aga	cca	ccc	agt	ttṭ	tat	atg	ggt	ctt	tac	tcc	ttc	ttt	gtg	ttt	gct	2832
Arg	Pro	Pro	Ser	Phe	Tyr	Met	Gly	Leu	Tyr	Ser	Phe	Phe	Val	Phe	Ala	
	930					935					940					
gct	ttc	ata	ttc	atg	aat	ggc	cag	ttc	acc	ata	ctt	tgc	gca	atg	ggt	2880
Ala	Phe	Ile	Phe	Met	Asn	Gly	Gln	Phe	Thr	Ile	Leu	Cys	Ala	Met	Gly	
945					950	•				955					960	
att	atg	gca	tcg	aaa	·tgg	tta	aat	ttg	agg	gct	gtg	aaa	aga	att	tta	2928
Ile	Met	Ala	Ser	Lys	Trp	Leu	Asn	Leu	Arg	Ala	Val	Lys	Arg	Ile	Leu	
				965					970					975		
cac	act	cca	atg	tca	tac	ata	gat	acc	aca	cct	ttg	gga	cgt	att	ctg	2976
His	Thr	Pro	Met	Ser	Tyr	Ile	Asp	Thr	Thr	Pro	Leu	Gly	Arg	Ile	Leu	
			980					985					990			
													-			
															gaa	3024
Asn	Arg	Phe	Thr	Lys	Asp	Thr	Asp	Ser	Leu	Asp	Asn	Glu	Leu	Thr	Glu	
	•	995				=	L000				1	L005				
															tgc	3072
Ser	Leu	Arg	Leu	Met	Thr	Ser	Gln	Phe	Ala	Asn	Ile	Val	Gly	Val	Cys	
:	1010				-	1015				-	L020					
					tac											3120
Val	Met	Cys	Ile	Val	Tyr	Leu	Pro	Trp	Phe	Ala	Ile	Ala	Ile	Pro	Phe	
102	5			;	1030				:	1035					1040	

							·									
				•					11							
ctt	ttg	gtc	atc	ttt	gtt	ctg	att	gct	gat	cat	tat	cag	agt	tct	ggt	3168
Leu	Leu	Val	Ile	Phe	Val	Leu	Ile	Ala	Asp	His	Tyr	Gln	Ser	Ser	Gly	
			:	1045				1	L050				1	.055		
апа	gaa	att	aaa	aga	ctt	gaa	gct	gtg	caa	cgg	tct	ttt	gtt	tac	aat	3216
				Arg												
ALG	GIG		1060					1065					1070			
		•	1000													
					.	~~+	~~~	ata	αat	aca	atc	aaa	gca	tac	cga	3264
															cga Arg	
Asn				Val	Leu			Mec	Азр	1111		-2- 1085		•		
		1075					1080				•					
											.		226	220	ata	3312
															atg	
Ser	Gln	Glu	Arg	, Phe	Leu	Ala	Lys	Ser	Asp			TTE	ASII	. шуз	ricc	
:	1090					1095					1100					
															.	
															att:	3360
Asn	Glu	ı Ala	a Gl	ү Туз	. Lei	ı Val	. Val	. Val	L Lev	ı Gln	Arg	Trp	val	. Gly	/ Ile	
110	5				1110	ס				1115	;				1120	
tto	ctt	gai	t at	g gti	t gci	t ato	gca	a tt	t gca	a cta	att	att	acg	g tta	a ttg	3408
															ı Leu	
				112					113					1139		
tat	: at	t ac	g aq	a qc	c tt	t cc	t at	t tc	c gc	g gct	t tca	a gti	t gg	t gt	t ttg	3456
															l Leu	
Cys			114					114					115			
			114						-							

ttg act tat gta tta caa ttg cct ggt cta tta aat acc att tta agg 3504

Leu	Thr	Tyr	Val	Leu	Gln	Leu	Pro	Gly	12 Leu	Asn	Thr	Ile	Leu	Arg
	1	155				1	1160			1	1165			

gca	atg	act	caa	aca	gag	aat	gac	atg	aat	agt	gcc	gaa	aga	ttg	gta	3552
Ala	Met	Thr	Gln	Thr	Glu	Asn	Asp	Met	Asn	Ser	Ala	Glu	Arg	Leu	Val	
1	170				J	175				1	L180					

aca	tat	gca	act	gaa	cta	cca	cta	gag	gca	tcc	tat	aga	aag	ccc	gaa	3600
Thr	Tyr	Ala	Thr	Glu	Leu	Pro	Leu	Glu	Ala	Ser	Tyr	Arg	Lys	Pro	Glu	
1189	5			1	L190				1	195				3	L200	

atg	aca	cct	cca	gag	tca	tgg	ccc	tca	atg	ggc	gaa	ata	att	ttt	gaa	3648		
Met	Thr	Pro	Pro	Glu	Ser	Trp	Pro	Ser	Met	Gly	Glu	Ile	Ile	Phe	Glu			
	1205						1210						1215					

- aat gtt gat ttt gcc tat aga cct ggt tta cct ata gtt tta aaa aat 3696
 Asn Val Asp Phe Ala Tyr Arg Pro Gly Leu Pro Ile Val Leu Lys Asn
 1220 1225 1230
- ctt aac ttg aat atc aag agt ggg gaa aaa att ggt atc tgt ggt cgt -3744

 Leu Asn Leu Asn Ile Lys Ser Gly Glu Lys Ile Gly Ile Cys Gly Arg

 1235 1240 1245
- aca ggt gct ggt aag tcc act att atg agt gcc ctt tac agg ttg aat 3792

 Thr Gly Ala Gly Lys Ser Thr Ile Met Ser Ala Leu Tyr Arg Leu Asn

 1250 1255 1260
- gaa ttg acc gca ggt aaa att tta att gac aat gtt gat ata agt cag 3840 Glu Leu Thr Ala Gly Lys Ile Leu Ile Asp Asn Val Asp Ile Ser Gln

13 1265 1270 1275 1280

ctg gga ctt ttc gat tta aga aga aaa tta gcc atc att cca caa gat 3888
Leu Gly Leu Phe Asp Leu Arg Arg Lys Leu Ala Ile Ile Pro Gln Asp
1285 1290 1295

cca gta tta ttt agg ggt acg att cgc aag aac tta gat cca ttt aat 3936
Pro Val Leu Phe Arg Gly Thr Ile Arg Lys Asn Leu Asp Pro Phe Asn
1300 1305 1310

gag cgt aca gat gac gaa tta tgg gat gca ttg gtg aga ggt ggt gct 3984
Glu Arg Thr Asp Asp Glu Leu Trp Asp Ala Leu Val Arg Gly Gly Ala
1315 1320 1325

atc gcc aag gat gac ttg ccg gaa gtg aaa ttg caa aaa cct gat gaa 4032

Ile Ala Lys Asp Asp Leu Pro Glu Val Lys Leu Gln Lys Pro Asp Glu

1330 1335 1340

aat ggt act cat ggt aaa atg cat aag ttc cat tta gat caa gca gtg 4080
Asn Gly Thr His Gly Lys Met His Lys Phe His Leu Asp Gln Ala Val

1345 1350 1355 1360

gaa gaa gag ggc tcc aat ttc tcc tta ggt gag aga caa cta tta gca 4128
Glu Glu Glu Gly Ser Asn Phe Ser Leu Gly Glu Arg Gln Leu Leu Ala

1365 1370 1375

tta aca agg gca ttg gtc cgc caa tca aaa ata ttg att ttg gat gag 4176
Leu Thr Arg Ala Leu Val Arg Gln Ser Lys Ile Leu Ile Leu Asp Glu
1380 1385 1390

14

gct	aca	tcc	tca	gtg	gac	tac	gaa	acg	gat	ggc	aaa	atc	caa	aca	cgt	4224
Ala	Thr	Ser	Ser	Val	Asp	Tyr	Glu	Thr	Asp	Gly	Lys	Ile	Gln	Thr	Arg	
	1	395				1	L 40 0]	L405				
											*					
att	gtt	gag	gaa	ttt	gga	gat	tgt	aca	atț	ttg	tgt	att	gct	cac	aga	4272
Ile	Val	Glu	Glu	Phe	Gly	Asp	Cys	Thr	Ile	Leu	Cys	Ile	Ala	His	Arg	
נ	410				3	1415		1420								
				gta												4320
Leu	Lys	Thr	Ile	Val	Asn	Tyr	Asp	Arg	Ile	Leu	Val	Leu	Glu	Lys	Gly	
1429	1425 1430								;	1435	1440					
																42.60
				ttc												4368
Glu	Val	Ala	Glu	Phe	Asp	Thr	Pro				Phe	Ser			Asp	
				1445					1450					1455		
													~~~	225	gat	4416
				agc												4410
Ser	Ile			Ser	Met	Cys				GTÀ	TTE		GIU 1470	ASII	wah	
			1460					1465					14/0			

ttc gag aac aga agt taa

4434

Phe Glu Asn Arg Ser

1475

<210> 2

<211> 1477

<212> PRT

15

<213> Saccharomyces cerevisiae

<400> 2

Met Thr Ile Thr Val Gly Asp Ala Val Ser Glu Thr Glu Leu Glu Asn

1 5 10 15

Lys Ser Gln Asn Val Val Leu Ser Pro Lys Ala Ser Ala Ser Ser Asp 20 25 30

Ile Ser Thr Asp Val Asp Lys Asp Thr Ser Ser Ser Trp Asp Asp Lys

35 40 45

Ser Leu Leu Pro Thr Gly Glu Tyr Ile Val Asp Arg Asn Lys Pro Gln
50 55 60

Thr Tyr Leu Asn Ser Asp Asp Ile Glu Lys Val Thr Glu Ser Asp Ile
65 70 75 80

Phe Pro Gln Lys Arg Leu Phe Ser Phe Leu His Ser Lys Lys Ile Pro

Glu Val Pro Gln Thr Asp Asp Glu Arg Lys Ile Tyr Pro Leu Phe His

Thr Asn Ile Ile Ser Asn Met Phe Phe Trp Trp Val Leu Pro Ile Leu
115 120 125

Arg Val Gly Tyr Lys Arg Thr Ile Gln Pro Asn Asp Leu Phe Lys Met

130 135 140

•

WO 99/10514

PCT/US98/17546

16

Asp Pro Arg Met Ser Ile Glu Thr Leu Tyr Asp Asp Phe Glu Lys Asn 145 150 155 160

Met Ile Tyr Tyr Phe Glu Lys Thr Arg Lys Lys Tyr Arg Lys Arg His

Pro Glu Ala Thr Glu Glu Glu Val Met Glu Asn Ala Lys Leu Pro Lys 180 185 190

His Thr Val Leu Arg Ala Leu Leu Phe Thr Phe Lys Lys Gln Tyr Phe
195 200 205

Met Ser Ile Val Phe Ala Ile Leu Ala Asn Cys Thr Ser Gly Phe Asn 210 215 220

Pro Met Ile Thr Lys Arg Leu Ile Glu Phe Val Glu Glu Lys Ala Ile
225 230 235 240

Phe His Ser Met His Val Asn Lys Gly Ile Gly Tyr Ala Ile Gly Ala
245 250 255

Cys Leu Met Met Phe Val Asn Gly Leu Thr Phe Asn His Phe Phe His

Thr Ser Gln Leu Thr Gly Val Gln Ala Lys Ser Ile Leu Thr Lys Ala 275 280 285

Ala Met Lys Lys Met Phe Asn Ala Ser Asn Tyr Ala Arg His Cys Phe

PCT/US98/17546 WO 99/10514

Pro Asn Gly Lys Val Thr Ser Phe Val Thr Thr Asp Leu Ala Arg Ile 

Glu Phe Ala Leu Ser Phe Gln Pro Phe Leu Ala Gly Phe Pro Ala Ile 

Leu Ala Ile Cys Ile Val Leu Leu Ile Val Asn Leu Gly Pro Ile Ala 

Leu Val Gly Ile Gly Ile Phe Phe Gly Gly Phe Phe Ile Ser Leu Phe 

Ala Phe Lys Leu Ile Leu Gly Phe Arg Ile Ala Ala Asn Ile Phe Thr 

Asp Ala Arg Val Thr Met Met Arg Glu Val Leu Asn Asn Ile Lys Met 

Ile Lys Tyr Tyr Thr Trp Glu Asp Ala Tyr Glu Lys Asn Ile Gln Asp 

Ile Arg Thr Lys Glu Ile Ser Lys Val Arg Lys Met Gln Leu Ser Arg 

Asn Phe Leu Ile Ala Met Ala Met Ser Leu Pro Ser Ile Ala Ser Leu 

Val Thr Phe Leu Ala Met Tyr Lys Val Asn Lys Gly Gly Arg Gln Pro
450 455 460

Gly Asn Ile Phe Ala Ser Leu Ser Leu Phe Gln Val Leu Ser Leu Gln
465 470 475 480

Met Phe Phe Leu Pro Ile Ala Ile Gly Thr Gly Ile Asp Met Ile Ile
485 490 495

Gly Leu Gly Arg Leu Gln Ser Leu Leu Glu Ala Pro Glu Asp Asp Pro 500 505 510

Asn Gln Met Ile Glu Met Lys Pro Ser Pro Gly Phe Asp Pro Lys Leu 515 520 525

Ala Leu Lys Met Thr His Cys Ser Phe Glu Trp Glu Asp Tyr Glu Leu 530 535 540

Asn Asp Ala Ile Glu Glu Ala Lys Gly Glu Ala Lys Asp Glu Gly Lys
545 550 555 560

Lys Asn Lys Lys Lys Arg Lys Asp Thr Trp Gly Lys Pro Ser Ala Ser 565 570 575

Thr Asn Lys Ala Lys Arg Leu Asp Asn Met Leu Lys Asp Arg Asp Gly
580 585 590

Pro Glu Asp Leu Glu Lys Thr Ser Phe Arg Gly Phe Lys Asp Leu Asn 595 600 605

WO 99/10514

19

Phe Asp Ile Lys Lys Gly Glu Phe Ile Met Ile Thr Gly Pro Ile Gly 610 615 620

Thr Gly Lys Ser Ser Leu Leu Asn Ala Met Ala Gly Ser Met Arg Lys 625 630 635 640

Thr Asp Gly Lys Val Glu Val Asn Gly Asp Leu Leu Met Cys Gly Tyr
645 650 655

Pro Trp Ile Gln Asn Ala Ser Val Arg Asp Asn Ile Ile Phe Gly Ser

Pro Phe Asn Lys Glu Lys Tyr Asp Glu Val Val Arg Val Cys Ser Leu 675 680 685

Lys Ala Asp Leu Asp Ile Leu Pro Ala Gly Asp Met Thr Glu Ile Gly 690 695 700

Glu Arg Gly Ile Thr Leu Ser Gly Gly Gln Lys Ala Arg Ile Asn Leu 705 710 715 720

Ala Arg Ser Val Tyr Lys Lys Lys Asp Ile Tyr Leu Phe Asp Asp Val

Leu Ser Ala Val Asp Ser Arg Val Gly Lys His Ile Met Asp Glu Cys
740 745 750

Leu Thr Gly Met Leu Ala Asn Lys Thr Arg Ile Leu Ala Thr His Gln

20

755 760 765

Leu Ser Leu Ile Glu Arg Ala Ser Arg Val Ile Val Leu Gly Thr Asp
770 780

Gly Gln Val Asp Ile Gly Thr Val Asp Glu Leu Lys Ala Arg Asn Gln
785 790 795 800

Thr Leu Ile Asn Leu Leu Gln Phe Ser Ser Gln Asn Ser Glu Lys Glu 805 810 815

Asp Glu Glu Glu Ala Val Val Ala Gly Glu Leu Gly Gln Leu Lys
820 825 830

Tyr Glu Ser Glu Val Lys Glu Leu Thr Glu Leu Lys Lys Lys Ala Thr

Glu Met Ser Gln Thr Ala Asn Ser Gly Lys Ile Val Ala Asp Gly His

Thr Ser Ser Lys Glu Glu Arg Ala Val Asn Ser Ile Ser Leu Lys Ile 865 870 875 880

Tyr Arg Glu Tyr Ile Lys Ala Ala Val Gly Lys Trp Gly Phe Ile Ala 885 890 895

Leu Pro Leu Tyr Ala Ile Leu Val Val Gly Thr Thr Phe Cys Ser Leu 900 905 910

PCT/US98/17546 WO 99/10514

Phe Ser Ser Val Trp Leu Ser Tyr Trp Thr Glu Asn Lys Phe Lys Asn

Arg Pro Pro Ser Phe Tyr Met Gly Leu Tyr Ser Phe Phe Val Phe Ala

Ala Phe Ile Phe Met Asn Gly Gln Phe Thr Ile Leu Cys Ala Met Gly

Ile Met Ala Ser Lys Trp Leu Asn Leu Arg Ala Val Lys Arg Ile Leu

. . . . . . . . . . . . . . .

His Thr Pro Met Ser Tyr Ile Asp Thr Thr Pro Leu Gly Arg Ile Leu

Asn Arg Phe Thr Lys Asp Thr Asp Ser Leu Asp Asn Glu Leu Thr Glu

Ser Lew Arg Lew Met Thr Ser Gln Phe Ala Asn Ile Val Gly Val Cys

Val Met Cys Ile Val Tyr Leu Pro Trp Phe Ala Ile Ala Ile Pro Phe

Leu Leu Val Ile Phe Val Leu Ile Ala Asp His Tyr Gln Ser Ser Gly

Arg Glu Ile Lys Arg Leu Glu Ala Val Gln Arg Ser Phe Val Tyr Asn

WO 99/10514

22

Asn Leu Asn Glu Val Leu Gly Gly Met Asp Thr Ile Lys Ala Tyr Arg

Ser Gln Glu Arg Phe Leu Ala Lys Ser Asp Phe Leu Ile Asn Lys Met 1090 1095 1100

Asn Glu Ala Gly Tyr Leu Val Val Val Leu Gln Arg Trp Val Gly Ile

105 1110 1115 1120

Phe Leu Asp Met Val Ala Ile Ala Phe Ala Leu Ile Ile Thr Leu Leu
1125 1130 1135

Cys Val Thr Arg Ala Phe Pro Ile Ser Ala Ala Ser Val Gly Val Leu 1140 1145 1150

Leu Thr Tyr Val Leu Gln Leu Pro Gly Leu Leu Asn Thr Ile Leu Arg

Ala Met Thr Gln Thr Glu Asn Asp Met Asn Ser Ala Glu Arg Leu Val

Thr Tyr Ala Thr Glu Leu Pro Leu Glu Ala Ser Tyr Arg Lys Pro Glu
185 1190 1195 1200

Met Thr Pro Pro Glu Ser Trp Pro Ser Met Gly Glu Ile Ile Phe Glu
1205 1210 1215

Asn Val Asp Phe Ala Tyr Arg Pro Gly Leu Pro Ile Val Leu Lys Asn

PCT/US98/17546 WO 99/10514

Leu Asn Leu Asn Ile Lys Ser Gly Glu Lys Ile Gly Ile Cys Gly Arg 

Thr Gly Ala Gly Lys Ser Thr Ile Met Ser Ala Leu Tyr Arg Leu Asn 

Glu Leu Thr Ala Gly Lys Ile Leu Ile Asp Asn Val Asp Ile Ser Gln 

Leu Gly Leu Phe Asp Leu Arg Arg Lys Leu Ala Ile Ile Pro Gln Asp 

Pro Val Leu Phe Arg Gly Thr Ile Arg Lys Asn Leu Asp Pro Phe Asn 

Glu Arg Thr Asp Asp Glu Leu Trp Asp Ala Leu Val Arg Gly Gly Ala 

Ile Ala Lys Asp Asp Leu Pro Glu Val Lys Leu Gln Lys Pro Asp Glu 

Asn Gly Thr His Gly Lys Met His Lys Phe His Leu Asp Gln Ala Val 

Glu Glu Glu Gly Ser Asn Phe Ser Leu Gly Glu Arg Gln Leu Leu Ala 

24

Leu Thr Arg Ala Leu Val Arg Gln Ser Lys Ile Leu Ile Leu Asp Glu

1380 1385 1390

Ala Thr Ser Ser Val Asp Tyr Glu Thr Asp Gly Lys Ile Gln Thr Arg 1395 1400 1405

Ile Val Glu Glu Phe Gly Asp Cys Thr Ile Leu Cys Ile Ala His Arg 1410 1415 1420

Leu Lys Thr Ile Val Asn Tyr Asp Arg Ile Leu Val Leu Glu Lys Gly
425 1430 1435 1440

Glu Val Ala Glu Phe Asp Thr Pro Trp Thr Leu Phe Ser Gln Glu Asp

Ser Ile Phe Arg Ser Met Cys Ser Arg Ser Gly Ile Val Glu Asn Asp 1460 1465 1470

Phe Glu Asn Arg Ser 1475

Inta ional Application No PCT/US 98/17546

		<u></u>	
A. CLASSIF IPC 6	FICATION OF SUBJECT MATTER C12N15/82 C12N15/31 A01H5/0	00	
According to	International Patent Classification (IPC) or to both national classif	cation and IPC	
B. FIELDS			
IPC 6	cumentation searched (classification system followed by classification C12N A01H		
Documentat	ion searched other than minimum documentation to the extent that	such documents are included in the fields sea	rched
Electronic d	ata base consulted during the international search (name of data l	pase and, where practical, search terms used)	
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the r	elevant passages	Relevant to claim No.
X	LU, YP., ET AL.: "AtMRP1 general Arabidopsis encodes a glutathio S-conjugate pump: Isolation and definition of a plant ATP-bindit transporter gene" PROCEEDINGS OF THE NATIONAL ACA SCIENCES OF USA., vol. 94, 22 July 1997, pages 82 XP002083621 WASHINGTON US see the whole document	ne functional ng cassette DEMY OF	1
P,X	WO 98 21938 A (UNIV PENNSYLVANI 28 May 1998 see page 90 - page 91; claims 1 figures 13-18		1
		-/	
		,	
X Furt	ther documents are listed in the continuation of box C.	X Patent family members are listed	in annex.
	ategories of cited documents :	"T" later document published after the inte	rnational filing date
"A" docum	ent defining the general state of the art which is not dered to be of particular relevance	or priority date and not in conflict with cited to understand the principle or the invention	the application but
filing		"X" document of particular relevance; the cannot be considered novel or cannot	t be considered to
which	ent which may throw doubts on priority claim(s) or his cited to establish the publication date of another on or other special reason (as specified)	involve an inventive step when the de "Y" document of particular relevance; the cannot be considered to involve an in	claimed invention
"O" docum	ment referring to an oral disclosure, use, exhibition or means	document is combined with one or m ments, such combination being obvious	ore other such docu-
"P" docum	ent published prior to the international filing date but than the priority date claimed	in the art.  "&" document member of the same paten	t family
Date of the	actual completion of theinternational search	Date of mailing of the international se	arch report
	November 1998	24/11/1998	
Name and	mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2	Authorized officer	
	NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Maddox, A	

3

Int ional Application No PCT/US 98/17546

Category 3	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
4	VOET, M, ET AL.: "S. cerevisiae chromosome VII reading frame ORF YGR281w" EMBL SEQUENCE ACCESSION NO. Z73066, 17 May 1996, XP002083622	1
4	KATZMANN, D.J., ET AL.: "Oligomycin resistance ATP-dependent permease YOR1" SWISS-PROT ACCESSION NO. P53049, 1 October 1996, XP002083623	1
A	see the whole document -& KATZMANN. D.J., ET AL.: "Expression of an ATP-binding cassette transporter-encoding gene (YOR1) is required for oligomycin resistance in Saccharomyces cerevisiae" MOLECULAR AND CELLULAR BIOLOGY, vol. 15, no. 12, December 1995, pages 6875-6883, XP002083629 see the whole document	1
Ą	WO 96 06175 A (PIONEER HI BRED INT) 29 February 1996 see the whole document	1-31
A	WO 96 20595 A (PROGUARD INC ;EMERSON RALPH W (US); CRANDALL BRADFORD G JR (US)) 11 July 1996 see the whole document	1-31
A	WO 95 06128 A (DEKALB GENETICS CORP) 2 March 1995 see page 37, line 25 - page 38, line 8	1-31
A	EP 0 644 262 A (TAKARA SHUZO CO) 22 March 1995 see sequence ID 9	1
A	ANZAI, H.,ET AL.: "Transgenic tobacco resistant to a bacterial disease by the detoxification of a pathogenic toxin" MOL. GEN. GENET., vol. 219, 1989, pages 492-494, XP002083624 see the whole document	1-31
А	DE LA FUENTE-MARTINEZ, J.M., ET AL.: "Expression of a bacterial phaseolotoxin resistant ornithyl transcarbamylase in transgenic tobacco confers resistance to Pesudomonas syringae pv. phaseolicola" BIOTECHNOLOGY, vol. 10, August 1992, pages 905-909, XP002083005 see the whole document	1-31
	-/	

3

Int. Ional Application No PCT/US 98/17546

		<u></u>
C.(Continu	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category 3	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 92 15685 A (RHONE POULENC AGROCHIMIE) 17 September 1992 see the whole document	1-31
A	WO 94 13790 A (SANOFI ELF ;ELF AQUITAINE (FR); PIGNARD ANNIE (FR); GREZES BESSET) 23 June 1994 see the whole document	1-31
A	WO 95 25114 A (UNIV TENNESSEE RES CORP) 21 September 1995 see page 16 - page 21	1-31
Α	WO 94 10303 A (UNIV KINGSTON) 11 May 1994 see the whole document	1-31
А	OW, D.W.: "Phytochelatin-mediated cadmium tolerance in Schizosaccharomyces pombe" IN VITRO CELL DEV BIOL, vol. 29p, October 1993, pages 213-219, XP002083625 see page 218, left-hand column, paragraph 2	1
Α	WO 91 06651 A (UNIV KINGSTON) 16 May 1991 see the whole document	1
A	WO 96 34959 A (AUCKLAND UNISERVICES LTD; GARDNER RICHARD CLAGUE (NZ); MACDIARMID) 7 November 1996 see the whole document	1

Information on patent family members

Int Ional Application No PCT/US 98/17546

		· · · · · · · · · · · · · · · · · · ·			<del></del>
Patent document cited in search rep		Publication date		atent family nember(s)	Publication date
WO 9821938	Α	28-05-1998	AU	5451198 A	10-06-1998
WO 9606175	Α	29-02-1996	US	5792931 A	11-08-1998
			AU	697831 B	15-10-1998
			AU	3491095 A	14-03-1996
			CA	2195784 A	29-02-1996
			EP	0775211 A	28-05-1997
			US	5716820 A	10-02-1998
			ZA	9506531 A	25-03-1996 
WO 9620595	Α	11-07-1996	AU	4744296 A	24-07-1996
			BR	9510180 A	14-10-1997
			CA	2208753 A	11-07-1996
			EP	0800345 A	15-10-1997
			PL	321179 A	24-11-1997
WO 9506128	Α	02-03-1995	AU	5640498 A	04-06-1998
			AU	684105 B	04-12-1997
			AU	7716994 A	21-03-1995
			BR	9407355 A	19-08-1997
			CA	2170260 A	02-03-1995
			EP	0721509 A	17-07-1996
			HU	74392 A	30-12-1996
			US	5780709 A	14-07-1998
			ZA	9406488 A	30-11-1995
EP 0644262	Α	22-03-1995	AU	677387 B	24-04-1997
			AU	6312994 A	01-12-1994
			CA	2124034 A	25-11-1994
			JP	7313172 A	05-12-1995
WO 9215685	Α	17-09-1992	FR	2673644 A	11-09-1992
			AU	660976 B	13-07-1995
			AU	1682092 A	06-10-1992
			BG	61275 B	30-04-1997
			BR	9204788 A	27-07-1993
			CA	2082042 A	06-09-1992
			CN	1065683 A	28-10-1992
			CZ	9203369 A	19-01-1994
			EP	0531498 A	17-03-1993

Information on patent family members

Inti onal Application No PCT/US 98/17546

	atent document in search report		Publication date		Patent family member(s)		Publication date
WO	9215685	Α		HU	65677	A	28-07-1994
				MX	9200917	Α	01-11-1992
				NZ	241829		27-04-1994
				PT	100203	Α	30-07-1993
WO	9413790	Α	23-06-1994	AU	5653294	_ <del>_</del>	04-07-1994
				CA	2151146	Α	23-06-1994
				EP	0672124	A	20-09-1995
WO	9525114		21-09-1995	US	5689039	Α	18-11-1997
				CA	2207851		21-09-1995
				EP	0767797	Α	16-04-1997
		•		JP	10500562	Т	20-01-1998
WO	9410303	Α	11-05-1994	AU	671160	В	15-08-1996
				AU	5173693	Α	24-05-1994
				AU	682140		18-09-1997
				AU	7177896	Α	06-02-1997
				EP	0666911	Α	16-08-1995
				JP	8504323	T	14-05-1996
				US	5489519		06-02-1996
				US	5766880	Α	16-06-1998
WO	9106651	Α	16-05-1991	US	5272085		21-12-1993
				AU	6542390		31-05-1991
				CA	2070395		01-05-1991
				EP	0494246		15-07-1992
				JP	5500900	T	25-02-1993
พก	9634959	Α	07-11-1996	AU	5517896	Α	21-11-1996